



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Analytic helicity amplitudes for two-loop five-gluon scattering: the single-minus case

Badger, Simon ; Brønnum-Hansen, Christian ; Hartanto, Heribertus Bayu ; Peraro, Tiziano

Abstract: We present a compact analytic expression for the leading colour two-loop five-gluon amplitude in Yang-Mills theory with a single negative helicity and four positive helicities. The analytic result is reconstructed from numerical evaluations over finite fields. The numerical method combines integrand reduction, integration-by-parts identities and Laurent expansion into a basis of pentagon functions to compute the coefficients directly from six-dimensional generalised unitarity cuts.

DOI: [https://doi.org/10.1007/jhep01\(2019\)186](https://doi.org/10.1007/jhep01(2019)186)

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-172597>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Badger, Simon; Brønnum-Hansen, Christian; Hartanto, Heribertus Bayu; Peraro, Tiziano (2019). Analytic helicity amplitudes for two-loop five-gluon scattering: the single-minus case. *Journal of High Energy Physics*, 2019(1):186.

DOI: [https://doi.org/10.1007/jhep01\(2019\)186](https://doi.org/10.1007/jhep01(2019)186)

Analytic helicity amplitudes for two-loop five-gluon scattering: the single-minus case

Simon Badger,^a Christian Brønnum-Hansen,^b Heribertus Bayu Hartanto^a
and Tiziano Peraro^c

^a*Institute for Particle Physics Phenomenology, Department of Physics, Durham University,
Durham DH1 3LE, United Kingdom*

^b*Higgs Centre for Theoretical Physics, School of Physics and Astronomy,
The University of Edinburgh,
Edinburgh EH9 3JZ, Scotland, U.K.*

^c*Physik-Institut, Universität Zürich,
Wintherturerstrasse 190, CH-8057 Zürich, Switzerland*

E-mail: simon.d.badger@durham.ac.uk, brønnum.hansen@ed.ac.uk,
heribertus.b.hartanto@durham.ac.uk, peraro@physik.uzh.ch

ABSTRACT: We present a compact analytic expression for the leading colour two-loop five-gluon amplitude in Yang-Mills theory with a single negative helicity and four positive helicities. The analytic result is reconstructed from numerical evaluations over finite fields. The numerical method combines integrand reduction, integration-by-parts identities and Laurent expansion into a basis of pentagon functions to compute the coefficients directly from six-dimensional generalised unitarity cuts.

KEYWORDS: Perturbative QCD, Scattering Amplitudes

ARXIV EPRINT: [1811.11699](https://arxiv.org/abs/1811.11699)

Contents

1	Introduction	1
2	Conventions and notation	2
3	Computational setup	3
3.1	Integrand parametrisation	4
3.2	Integration-by-parts identities	6
3.3	Map to pentagon functions	7
3.4	Laurent expansion	8
4	Results	9
5	Conclusions	12

1 Introduction

The increasing need for high precision predictions for Standard Model processes at Hadron colliders has set out a priority list of new perturbative calculations required to keep theoretical uncertainties in line with experimental errors [1]. A large category of these processes requires unknown two-loop $2 \rightarrow 3$ scattering amplitudes. Owing to the high degree of difficulty of these calculations, recent years have seen an increasing effort to find new techniques capable of providing these results.

Due to the success of automated algorithms for the numerical computation of one-loop amplitudes, there has been considerable interest in extending the established methods of integrand reduction [2–6] and (generalised) unitarity [7–14] to two loops and beyond [15–26]. For planar $2 \rightarrow 3$ scattering in Quantum Chromodynamics (QCD) this effort has led to analytic results for the all-plus helicity two-loop amplitude [27–33]. The remaining helicity configurations have been obtained numerically [34–37]. Some groups have been able to construct solutions to the integration-by-parts reduction identities analytically [38, 39] yet no complete amplitudes were obtained in compact form.

One of the major challenges in this program has been to understand how to efficiently build in simplifications from integration-by-parts identities (IBPs) [40, 41] that first appear at two loops [42–48]. A conventional approach to solving this reduction problem with the Laporta algorithm [49] can be extremely computationally intensive, especially in cases with many kinematic scales. On-going work continues to produce more and more efficient algorithms [50–52]. The use of finite field arithmetic has also been shown to provide a highly efficient method which can avoid traditional bottlenecks [53]. It is this last approach which we build on in this paper. It has also been demonstrated how this technique can be applied to compute scattering amplitudes through multivariate functional reconstruction [25].

Another major step in the evaluation of five-point two-loop scattering amplitudes is the computation of a basis of integral functions. Considerable progress on this front has been made recently and the analytic evaluation of many of the two loop integrals required after reduction has been completed with the help of differential equation methods [54–65].

Analytic results can offer many benefits over numerical algorithms. The one-loop amplitudes for five-gluon scattering, first derived in 1993 by Bern, Dixon and Kosower [66], are strikingly simple. One immediate consequence of this is that amplitudes are fast and stable to evaluate numerically and well suited for Monte Carlo integration. Analytic results also give us more insight into the structure of on-shell amplitudes in gauge theory. Simplicity in maximally super-symmetric Yang-Mills theory has enabled huge leaps into the structure of perturbative amplitudes based on constraints from universal behaviour in physical limits [67–70]. While in QCD these constraints are not quite enough to fix the amplitudes (such techniques have been applied in the computation of the QCD soft anomalous dimension [71]), it would be an extremely powerful tool if the function space of multi-loop amplitudes could be better understood in general gauge theories.

In this paper we present new, analytic results for the scattering of five gluons in pure Yang-Mills at two loops in which one gluon has negative helicity and the remaining gluons have positive helicities. We employ finite field numerics to a combined system of integrand reduction, integration-by-parts identities and expansion into a basis of pentagon functions. After multiple evaluations we were able to reconstruct the analytic form of the amplitude.

We outline our conventions and notation in section 2. We then describe the numerical algorithm used to map the coefficients of a pentagon function basis for the finite remainder of the two-loop amplitude from generalised unitarity cuts with six-dimensional tree amplitudes in section 3. This numerical algorithm is then sampled using finite field arithmetic and the rational coefficients of the polylogarithmic pentagon functions are reconstructed as functions of momentum twistor variables. We present our results in section 4 before drawing some brief conclusions.

2 Conventions and notation

We compute the leading colour contribution to five-gluon scattering in pure Yang-Mills in the fundamental trace basis:

$$\mathcal{A}^{(L)}(1_g, 2_g, 3_g, 4_g, 5_g) = n^L g_s^3 \sum_{\sigma \in S_5/Z_5} \text{tr}(T^{a_{\sigma(1)}} T^{a_{\sigma(2)}} T^{a_{\sigma(3)}} T^{a_{\sigma(4)}} T^{a_{\sigma(5)}}) \times A^{(L)}(\sigma(1)_g, \sigma(2)_g, \sigma(3)_g, \sigma(4)_g, \sigma(5)_g) \quad (2.1)$$

where L is the number of loops and we have extracted a normalisation defined by,

$$n = m_\epsilon N_c \alpha_s / (4\pi), \quad \alpha_s = g_s^2 / (4\pi), \quad m_\epsilon = i(4\pi)^\epsilon e^{-\epsilon\gamma_E}. \quad (2.2)$$

We further expand the amplitudes around $d_s = 2$ where $d_s = g^\mu{}_\mu$ is the spin dimension,

$$A^{(1)}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+) = (d_s - 2) A^{(1),[1]}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+), \quad (2.3a)$$

$$A^{(2)}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+) = \sum_{i=0}^2 (d_s - 2)^i A^{(2),[i]}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+). \quad (2.3b)$$

This is useful since the $d_s = 2$ limit behaves like a supersymmetric amplitude¹ where additional cancellations and simplifications can be seen. In the case of the single-minus helicity configuration, it was already observed that $A^{(2),[0]}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+) = \mathcal{O}(\epsilon)$ [34].

Since the tree-level helicity amplitude is zero, the universal infrared (IR) poles take a very simple form [72–75],

$$A^{(1),[1]}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+) = F^{(1),[1]}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+) + \mathcal{O}(\epsilon), \quad (2.4a)$$

$$A^{(2),[1]}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+) = \left[-\frac{r_\Gamma}{\epsilon^2} \sum_{i=1}^5 \left(\frac{\mu_R^2 e^{\gamma_E}}{-s_{i,i+1}} \right)^\epsilon \right] A^{(1),[1]}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+) + F^{(2),[1]}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+) + \mathcal{O}(\epsilon), \quad (2.4b)$$

$$A^{(2),[2]}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+) = F^{(2),[2]}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+) + \mathcal{O}(\epsilon), \quad (2.4c)$$

where

$$r_\Gamma = \frac{\Gamma^2(1-\epsilon)\Gamma(1+\epsilon)}{\Gamma(1-2\epsilon)}. \quad (2.5)$$

Note that, in eq. (2.4b), the one-loop amplitude² $A^{(1),[1]}$ needs to be expanded up to $\mathcal{O}(\epsilon^2)$. In this paper we will present a direct computation of the finite remainder F .

3 Computational setup

The kinematic parts of the amplitude are written using a momentum twistor [76] parametrisation, as described in previous works [27, 34]. We decompose the amplitude into an integrand basis, using the method of integrand reduction via generalised unitarity. We then reduce the amplitude to master integrals by solving IBPs. The master integrals are in turn expressed as combinations of known pentagon functions, using the expressions computed in reference [63].

The algorithm is implemented numerically over finite fields. The Laurent expansion in ϵ of the results is obtained by performing a full reconstruction of its dependence on the dimensional regulator ϵ , for fixed numerical values of the kinematic variables. The Laurent expansion of the reconstructed function of ϵ thus provides a numerical evaluation of the ϵ -expansion of the final result. Finally, the full dependence of the expanded result on the kinematic variables is reconstructed from multiple numerical evaluations, using a modified version of the multi-variate reconstruction techniques presented in reference [25].

In this section we provide more details on our computational setup and the various steps of the calculation outlined above.

¹One can show that setting $d_s = 2$ is equivalent to performing a supersymmetric decomposition [66] with $n_f = 0$ adjoint fermions and $n_s = -1$ (complex) adjoint scalars, and that this yields a linear combination of supersymmetric contributions.

²Expressions for the one-loop amplitudes can be obtained from the authors on request.




Figure 1. Two-loop five-point master topologies. All external momenta are considered out-going, arrows indicate loop momenta directions.

3.1 Integrand parametrisation

We define an integral family by a complete, minimal set of propagators and irreducible scalar products (ISPs):

$$\begin{aligned}
 G_{a_1a_2a_3a_4a_5a_6a_7a_8a_9a_{10}a_{11}} &= \int \frac{d^d k_1}{i\pi^{d/2}e^{-\epsilon\gamma_E}} \frac{d^d k_2}{i\pi^{d/2}e^{-\epsilon\gamma_E}} \\
 &\times \frac{1}{k_1^{2a_1}} \frac{1}{(k_1 - p_1)^{2a_2}} \frac{1}{(k_1 - p_1 - p_2)^{2a_3}} \frac{1}{(k_1 + p_4 + p_5)^{2a_4}} \\
 &\times \frac{1}{k_2^{2a_5}} \frac{1}{(k_2 - p_5)^{2a_6}} \frac{1}{(k_2 - p_4 - p_5)^{2a_7}} \frac{1}{(k_1 + k_2)^{2a_8}} \\
 &\times \frac{1}{(k_1 + p_5)^{2a_9}} \frac{1}{(k_2 + p_1)^{2a_{10}}} \frac{1}{(k_2 + p_1 + p_2)^{2a_{11}}}, \quad (3.1)
 \end{aligned}$$

where the exponents, a_i , are integers and $d = 4 - 2\epsilon$. The three master topologies, shown in figure 1, are

$$\text{Pentabox:} \quad G_{111111111a_9a_{10}a_{11}} \quad (3.2)$$


$$\text{Hexatriangle:} \quad G_{1111111a_711a_{10}a_{11}} \quad (3.3)$$

$$\text{Heptabubble:} \quad G_{211111a_6a_711a_{10}a_{11}} \quad (3.4)$$

while propagators with unspecified exponents, a_j , correspond to ISPs (i.e. $a_j \leq 0$).


All lower point topologies are obtained by systematically pinching the propagators of the master topologies. Topologies with scaleless integrals are discarded since we work in dimensional regularisation. Pinching of propagators from different master topologies can lead to the same sub-topology. This happens in particular when all five cyclic permutations of the external momenta are included. In these cases the assignment to a master topology is not unique. The full set of 57 distinct topologies with a specific choice of master topology assignment is shown in figure 2.

We parametrise the integrand numerators by writing the most general polynomials in the ISPs subject to a power counting constraint from renormalizability considerations. As



(a) Topologies associated with the pentabox master topology (top left), see (3.2).

(b) Topologies associated with the hexatriangle master topology (top left), see (3.3).



(c) Topologies associated with the heptabubble master topology (top left), see (3.4).

(d) Topologies with divergent cuts that must be computed simultaneously with sub-topologies of the heptabubble master topology in figure 2c.

Figure 2. All distinct two-loop five-point topologies.

an example, the pentabox of figure 1a has the numerator parametrisation,

$$\Delta \left(\begin{array}{c} 5 \\ \diagdown \quad \diagup \\ \text{---} \quad \text{---} \\ \diagup \quad \diagdown \\ 4 \end{array} \begin{array}{c} \xleftarrow{k_2} \quad \xrightarrow{k_1} \\ \diagup \quad \diagdown \\ \text{---} \quad \text{---} \\ \diagdown \quad \diagup \\ 3 \end{array} \begin{array}{c} 1 \\ \diagdown \quad \diagup \\ \text{---} \quad \text{---} \\ \diagup \quad \diagdown \\ 3 \end{array} \right) = \sum c_{(1,1,1,1,1,1,1,1,a_9,a_{10},a_{11})} (k_1+p_5)^{-2a_9} (k_2+p_1)^{-2a_{10}} (k_2+p_1+p_2)^{-2a_{11}} \quad (3.5)$$

where the sum is truncated by the constraints on the exponents:

$$-5 \leq a_9 \leq 0, \quad (3.6)$$

$$-4 \leq a_{10} + a_{11} \leq 0, \quad (3.7)$$

$$-7 \leq a_9 + a_{10} + a_{10} \leq 0. \quad (3.8)$$

Each topology has $11 - n$ ISPs where n is the number of distinct propagators. The five cyclic permutations of the external legs give a total of 425 irreducible numerators.

Integrand representations of the form (3.5) are less compact than representations making use of, for example, local integrands, spurious integrands, and extra-dimensional ISPs [27, 28, 32, 34, 77–79]. However, in our set-up the integrand is only sampled numerically and not analytically reconstructed. Simplification at the integrand level is therefore not a priority. Because our final integrated amplitude does not depend on the choice of integrand parametrisation, we have chosen a form which is directly compatible with IBPs, rather than one yielding a compact integrand representation. On the other hand there is potential for considerable improvements in the efficiency of the algorithm if a simpler integrand form could be identified.

We take a top down approach to solving the complete system of integrands which, apart from the basis choice described above, is identical to the approach taken in ref. [34]. The tree amplitudes used to compute the generalised unitarity cuts are evaluated by contracting Berends-Giele currents [80] as described in [25] and the six-dimensional spinor-helicity formalism [81]. Eight topologies, shown in figure 2d, have divergent cuts and their integrand coefficients are determined simultaneously with sub-topologies in the heptabubble group (see figure 2c). This follows the approach used previously in references [26, 34].

The numerical sampling of the integrand can show quickly which coefficients vanish and hence what integrals require further reduction using IBPs. The number of non-vanishing coefficients at the integrand level split into the components of $d_s = 2, (d_s - 2), (d_s - 2)^2$ are 4387, 14565, 4420 respectively. We find the maximum rank to be 5 for genuine two-loop topologies and rank 6 for a few integrals in the $(d_s - 2)^2$ component of the amplitude that can be written as (1-loop)² integrals, see figure 2a.

At the end of the integrand reduction stage, the colour ordered amplitude can be written as

$$A^{(2)}(1, 2, 3, 4, 5) = \sum_{\mathbf{a}} c_{\mathbf{a}}^{[\partial A]} G_{\mathbf{a}}, \quad (3.9)$$

where we sum over the tuples $\mathbf{a} = (a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9, a_{10}, a_{11})$. The coefficients $c_{\mathbf{a}}^{[\partial A]}$ are rational functions in the momentum twistor variables only.

3.2 Integration-by-parts identities

Each integral appearing in (3.9) is reduced to a set of master integrals J_k ,

$$G_{\mathbf{a}} = \sum_k c_{\mathbf{a}k}^{[\text{IBP}]} J_k, \quad (3.10)$$

where the sum runs over 155 master integrals (remembering that we include the 5 cyclic permutations of the integral family G). The reduction is obtained by solving a traditional Laporta system of IBP equations [49]. The system is generated in MATHEMATICA with the help of LITERED [82], and solved over finite fields, for numerical values of ϵ and the kinematic invariants, with a custom general-purpose linear solver for sparse systems of equations. The master integrals are chosen to be the uniform weight functions identified

by Gehrmann, Henn and Lo Presti [63]. The $c_{\mathbf{a}k}^{[\text{IBP}]}$ are rational functions in the momentum twistor variables and the dimensional regularisation parameter ϵ .

3.3 Map to pentagon functions

Our next step is to expand the master integrals into a basis of pentagon functions defined by Gehrmann, Henn and Lo Presti. These functions can be written in terms of Goncharov Polylogarithms. We take the results of expanding the master integrals in ϵ from reference [63],

$$J_k = \sum_{x=0}^4 \sum_l c_{kl;x}^{[f]} \epsilon^x m_{l;x}(f) + \mathcal{O}(\epsilon^5), \quad (3.11)$$

where $m_{l;x}(f)$ are monomials in the pentagon functions (note that the coefficients $c_{kl;x}^{[f]}$ depend on the choice of the pentagon functions f).

The amplitude can thus be written as a combination of pentagon functions

$$A^{(2)}(1, 2, 3, 4, 5) = \sum_{l,x} c_{l;x}^{[A]} m_{l;x}(f) + \mathcal{O}(\epsilon), \quad (3.12)$$

where the coefficients are defined through matrix multiplication, from the three reduction steps

$$c_{l;x}^{[A]} = \sum_{\mathbf{a},k} \sum_{x=0}^4 c_{\mathbf{a}}^{[\partial A]} c_{\mathbf{a}k}^{[\text{IBP}]} c_{kl;x}^{[f]} \epsilon^x. \quad (3.13)$$

We recall that, in the previous equation, there is also an implicit dependence on ϵ coming from the $c_{\mathbf{a}k}^{[\text{IBP}]}$, which were defined in (3.10) to be the full coefficients of the IBP reduction. Hence, the coefficients $c_{l;x}^{[A]}$ are rational functions of ϵ which need to be expanded, as we will explain in the next subsection.

The final step of the algorithm is to perform the same decomposition for the universal IR poles in (2.4b). For this we need the one-loop master integrals expanded up to weight four and written in the same alphabet as the two-loop integrals. These results were obtained directly from the differential equations in a canonical basis.³ We then write the poles analytically as

$$\left[-\frac{r_\Gamma}{\epsilon^2} \sum_{j=1}^5 \left(\frac{\mu_R^2 e^{\gamma_E}}{-s_{j,j+1}} \right)^\epsilon \right] A^{(1),[i]}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+) = \sum_{l,x} c_{l;x}^{[\text{IR}]} m_{l;x}(f) + \mathcal{O}(\epsilon). \quad (3.14)$$

Our numerical algorithm can then compute the difference:

$$c_{l;x}^{[F]} = c_{l;x}^{[A]} - c_{l;x}^{[\text{IR}]} \quad (3.15)$$

which we will expand in ϵ to find the finite remainder. At this point we have constructed a numerical algorithm which combines integrand reduction, IBP reduction and expansion of the master integrals into a basis of polylogarithms. This algorithm can be used to compute the finite remainder of the two-loop amplitude through evaluations of generalised unitarity cuts over finite fields.

³We are very grateful to Adriano Lo Presti for assistance in setting up the differential equations used in [59].

3.4 Laurent expansion

In the previous subsections we described a numerical calculation over finite fields of the coefficients $c_{l;x}^{[F]}$. They are used in order to write the finite remainder, $F^{(2),[i]}$, of the amplitude in terms of known pentagon functions. The coefficients, computed as described above, are rational functions of ϵ . However, because the calculation uses the expansion in (3.11) for the master integrals in terms of pentagon functions, it is only valid up to $\mathcal{O}(\epsilon)$. Here we are interested in the finite part of the Laurent expansion in ϵ .

As mentioned before, in order to obtain this Laurent expansion, we first perform a full reconstruction of the functions $c_{l;x}^{[F]}$ in ϵ , for numerical values over finite fields of the momentum twistor variables. The reconstructed function can thus be expanded in ϵ up to the desired order. This yields a decomposition of the form

$$c_{l;x}^{[F]} = \sum_{y=-4}^0 c_{l;x,y}^{[F]} \epsilon^y + \mathcal{O}(\epsilon), \quad (3.16)$$

where we are interested in the finite parts $c_{l;x,0}^{[F]}$, while $c_{l;x,y}^{[F]} = 0$ for $y < 0$. The finite remainder is therefore

$$F^{(2),[1]}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+) = \sum_l \sum_{x=0}^4 c_{l;x,0}^{[F]} m_{l;x}(f) + \mathcal{O}(\epsilon), \quad (3.17)$$

with $c_{l;x,0}^{[F]}$ defined by the Laurent expansion in (3.16).

The algorithm described above numerically computes the coefficients $c_{l;x,0}^{[F]}$ of the finite remainder of the amplitude over finite fields, for any numerical value of the kinematic invariants represented by the momentum twistor variables. Full analytic formulas for the coefficients $c_{l;x,0}^{[F]}$, as rational functions of the momentum twistor variables, are reconstructed from multiple numerical evaluations. The number of sample points for the three components $d_s = 2, (d_s - 2), (d_s - 2)^2$ are 3, 2214, 22886 respectively and sampling over one finite field is sufficient. For this purpose, we use a slightly improved version of the multivariate reconstruction techniques presented in reference [25].⁴ We note the large difference in the number of sample points needed for the different components in the $d_s = 2$ expansion. This happens since the coefficients of the pentagon function basis used in the fit contains higher powers of spurious poles for $(d_s - 2)^2$ than for $(d_s - 2)$ even though the integrals and topologies appearing are much simpler. Once these expressions are collected and written in terms of the finite integral functions described in the next section, both amplitudes take similarly compact forms. The difference in time to perform the fit was not prohibitive in this case so further optimisation of the basis before the fit was unnecessary.

In the next section we give a compact form of this result, obtained from the one in terms of momentum twistor variables, after converting it into spinor products and momentum invariants via some additional algebraic manipulations.

⁴These improvements concern performance, memory usage, and parallelization, and will be described in a later publication.

4 Results

We present a compact form of the amplitude by making use of the symmetry $(1, 2, 3, 4, 5) \rightarrow (1, 5, 4, 3, 2)$ and extracting an overall phase written in terms of spinor products,

$$F^{(L),[i]}(1_g^-, 2_g^+, 3_g^+, 4_g^+, 5_g^+) = \frac{[25]^2}{[12]\langle 23\rangle\langle 34\rangle\langle 45\rangle[51]} \left(F_{\text{sym}}^{(L),[i]}(1, 2, 3, 4, 5) + F_{\text{sym}}^{(L),[i]}(1, 5, 4, 3, 2) \right), \quad (4.1)$$

where L labels the loop order and i labels the component in the expansion around $d_s = 2$. The known result at one-loop can be written as:

$$F_{\text{sym}}^{(1),[1]}(1, 2, 3, 4, 5) = \frac{\text{tr}_+(2315)^2 \text{tr}_+(1243)}{3s_{25}^2 s_{23} s_{34} s_{15}} - \frac{\text{tr}_+(2543)}{6s_{34}}, \quad (4.2)$$

where $\text{tr}_+(ijkl) = \frac{1}{2} \text{tr}((1 + \gamma_5)\not{p}_i \not{p}_j \not{p}_k \not{p}_l)$ and $s_{ij} = (p_i + p_j)^2$.

The finite parts of the two-loop amplitude can be written compactly in terms of weight two functions, just as at one-loop. We therefore follow the same strategy as at one-loop to find a basis of integral functions free of large cancellations due to spurious singularities. We find that a convenient basis for the $d_s - 2$ component of the amplitude is

$$\begin{aligned} F_{\text{sym}}^{(2),[1]}(1, 2, 3, 4, 5) = & c_{51}^{(2)} F_{\text{box}}^{(2)}(s_{23}, s_{34}, s_{15}) + c_{51}^{(1)} F_{\text{box}}^{(1)}(s_{23}, s_{34}, s_{15}) + c_{51}^{(0)} F_{\text{box}}^{(0)}(s_{23}, s_{34}, s_{15}) \\ & + c_{34}^{(2)} F_{\text{box}}^{(2)}(s_{12}, s_{15}, s_{34}) + c_{34}^{(1)} F_{\text{box}}^{(1)}(s_{12}, s_{15}, s_{34}) + c_{34}^{(0)} F_{\text{box}}^{(0)}(s_{12}, s_{15}, s_{34}) \\ & + c_{45} F_{\text{box}}^{(0)}(s_{12}, s_{23}, s_{45}) + c_{34;51} \hat{L}_1(s_{34}, s_{15}) + c_{51;23} \hat{L}_1(s_{15}, s_{23}) + c_{\text{rat}}, \end{aligned} \quad (4.3)$$

and

$$\begin{aligned} F_{\text{sym}}^{(2),[2]}(1, 2, 3, 4, 5) = & d_{34}^{(3)} F_{\text{box}}^{(3)}(s_{12}, s_{15}, s_{34}) + d_{34}^{(2)} F_{\text{box}}^{(2)}(s_{12}, s_{15}, s_{34}) \\ & + d_{34;51}^{(3)} \hat{L}_3(s_{34}, s_{15}) + d_{34;51}^{(2)} \hat{L}_2(s_{34}, s_{15}) \\ & + d_{51;23}^{(3)} \hat{L}_3(s_{15}, s_{23}) + d_{51;23}^{(2)} \hat{L}_2(s_{15}, s_{23}) + d_{\text{rat}}, \end{aligned} \quad (4.4)$$

for the $(d_s - 2)^2$ amplitude.

The integral functions are written in terms of simple logarithms and di-logarithms. All weight one functions appear as logarithms of ratios of kinematic invariants,

$$L_k(s, t) = \frac{\log(t/s)}{(s - t)^k}, \quad (4.5)$$

where the singular behaviour is removed by defining,

$$\hat{L}_0(s, t) = L_0(s, t), \quad (4.6a)$$

$$\hat{L}_1(s, t) = L_1(s, t), \quad (4.6b)$$

$$\hat{L}_2(s, t) = L_2(s, t) + \frac{1}{2(s - t)} \left(\frac{1}{s} + \frac{1}{t} \right), \quad (4.6c)$$

$$\hat{L}_3(s, t) = L_3(s, t) + \frac{1}{2(s - t)^2} \left(\frac{1}{s} + \frac{1}{t} \right). \quad (4.6d)$$

At weight two all functions can be written in terms of the six-dimensional box function,

$$F_{\text{box}}^{(-1)}(s, t, m^2) = \text{Li}_2\left(1 - \frac{s}{m^2}\right) + \text{Li}_2\left(1 - \frac{t}{m^2}\right) + \log\left(\frac{s}{m^2}\right) + \log\left(\frac{t}{m^2}\right) - \frac{\pi^2}{6}, \quad (4.7a)$$

$$F_{\text{box}}^{(0)}(s, t, m^2) = \frac{1}{u(s, t, m^2)} F_{\text{box}}^{(-1)}(s, t, m^2), \quad (4.7b)$$

$$F_{\text{box}}^{(1)}(s, t, m^2) = \frac{1}{u(s, t, m^2)} \left[F_{\text{box}}^{(0)}(s, t, m^2) + \hat{L}_1(s, m^2) + \hat{L}_1(m^2, t) \right], \quad (4.7c)$$

$$F_{\text{box}}^{(2)}(s, t, m^2) = \frac{1}{u(s, t, m^2)} \left[F_{\text{box}}^{(1)}(s, t, m^2) + \frac{s - m^2}{2t} \hat{L}_2(s, m^2) + \frac{m^2 - t}{2s} \hat{L}_2(m^2, t) - \left(\frac{1}{s} + \frac{1}{t}\right) \frac{1}{4m^2} \right], \quad (4.7d)$$

$$F_{\text{box}}^{(3)}(s, t, m^2) = \frac{1}{u(s, t, m^2)} \left[F_{\text{box}}^{(2)}(s, t, m^2) - \frac{(s - m^2)^2}{6t^2} \hat{L}_3(s, m^2) - \frac{(m^2 - t)^2}{6s^2} \hat{L}_3(m^2, t) - \left(\frac{1}{s} + \frac{1}{t}\right) \frac{1}{6m^4} \right], \quad (4.7e)$$

where $u(s, t, m^2) = m^2 - s - t$.

These functions serve the same purpose as the L s and L functions introduced by Bern, Dixon, and Kosower in [66, 83]. The $\hat{L}_i(s, t)$ are finite as $s \rightarrow t$ and the $F_{\text{box}}^{(i)}(s, t, m^2)$ are finite as $s \rightarrow -t + m^2$. The definitions have been changed very slightly with respect to the L s and L functions since the singularities from the box functions at $m^2 - s - t$ have been removed without introducing additional singularities in $s - m^2$ or $t - m^2$.

For the $(d_s - 2)$ amplitude the coefficients are:

$$c_{51}^{(2)} = \frac{5s_{23}s_{34} \text{tr}_+(1234)^2 \text{tr}_+(1542)^2}{s_{12}s_{15} \text{tr}_+(2543)^2}, \quad (4.8a)$$

$$c_{51}^{(1)} = -\frac{\text{tr}_+(1234)^2 \text{tr}_+(1534) \text{tr}_+(2453)^2}{6s_{12}s_{34}s_{35} \text{tr}_+(2543)^2}, \quad (4.8b)$$

$$\begin{aligned} c_{51}^{(0)} = & \frac{s_{15}s_{45} \text{tr}_+(1234)}{3 \text{tr}_+(2543)} - \frac{s_{15}s_{24}s_{45} \text{tr}_+(1234)^2}{6s_{12} \text{tr}_+(2543)^2} - \frac{\text{tr}_+(1234)^2 \text{tr}_+(1542)}{6 \text{tr}_+(2543)^2} \\ & - \frac{s_{23}s_{24} \text{tr}_+(1234) \text{tr}_+(1543)}{6 \text{tr}_+(2543)^2} - \frac{s_{12}s_{23}s_{34} \text{tr}_+(1542) \text{tr}_+(1543)}{3s_{15} \text{tr}_+(2543)^2} \\ & + \frac{2s_{23} \text{tr}_+(1234) \text{tr}_+(1543)^2}{3s_{15} \text{tr}_+(2543)^2} - \frac{s_{24} \text{tr}_+(1234) \text{tr}_+(1543)^2}{6s_{15} \text{tr}_+(2543)^2} \\ & - \frac{s_{24} \text{tr}_+(1234)^2 \text{tr}_+(1543)^2}{6s_{12}s_{15}s_{34} \text{tr}_+(2543)^2} - \frac{s_{24} \text{tr}_+(1235) \text{tr}_+(1243) \text{tr}_+(1543)^2}{6s_{12}s_{15}s_{34} \text{tr}_+(2543)^2} \\ & - \frac{\text{tr}_+(1234) \text{tr}_+(1543) \text{tr}_+(2453)}{2 \text{tr}_+(2543)^2}, \end{aligned} \quad (4.8c)$$

$$c_{34}^{(2)} = \frac{5}{2} s_{12}^2 s_{15}^2, \quad (4.9a)$$

$$c_{34}^{(1)} = \frac{\text{tr}_+(1245) \text{tr}_+(1534) \text{tr}_+(1543)}{3s_{15}s_{34}s_{45}} + \frac{1}{12} s_{12}s_{15}s_{34}, \quad (4.9b)$$

$$\begin{aligned}
 c_{34}^{(0)} = & -\frac{s_{15}s_{23} \operatorname{tr}_+(1234)}{12 \operatorname{tr}_+(2543)} - \frac{\operatorname{tr}_+(1234)^2 \operatorname{tr}_+(1532)}{12s_{12}s_{34} \operatorname{tr}_+(2543)} + \frac{s_{25} \operatorname{tr}_+(1234) \operatorname{tr}_+(1543)}{12s_{34} \operatorname{tr}_+(2543)} \\
 & + \frac{s_{12} \operatorname{tr}_+(1532) \operatorname{tr}_+(1543)}{12s_{15} \operatorname{tr}_+(2543)} + \frac{s_{23}s_{25} \operatorname{tr}_+(1234) \operatorname{tr}_+(1543)^2}{12s_{15}s_{34} \operatorname{tr}_+(2543)^2} \\
 & + \frac{s_{12} \operatorname{tr}_+(1543)^2 \operatorname{tr}_+(2354)}{12s_{15}s_{34}s_{45} \operatorname{tr}_+(2543)} + \frac{s_{25} \operatorname{tr}_+(2543)}{12s_{34}} - \frac{1}{3}s_{12}s_{15},
 \end{aligned} \tag{4.9c}$$

$$c_{45} = -\frac{s_{13}s_{45} \operatorname{tr}_+(1234)^2 \operatorname{tr}_+(1534)}{6s_{12}s_{34} \operatorname{tr}_+(2543)^2} + \frac{s_{23}^3 s_{34} \operatorname{tr}_+(1543)}{6 \operatorname{tr}_+(2543)^2} - \frac{s_{13}s_{23} \operatorname{tr}_+(1243) \operatorname{tr}_+(1543)^2}{6s_{15}s_{34} \operatorname{tr}_+(2543)^2}, \tag{4.10}$$

$$\begin{aligned}
 c_{34;51} = & \frac{s_{12}s_{15}s_{34}s_{45}}{6 \operatorname{tr}_+(2543)} - \frac{11 \operatorname{tr}_+(1234) \operatorname{tr}_+(1543)}{6 \operatorname{tr}_+(2543)} + \frac{\operatorname{tr}_+(1234)^2 \operatorname{tr}_+(1542) \operatorname{tr}_+(1543)}{6s_{12}s_{34} \operatorname{tr}_+(2543)^2} \\
 & - \frac{s_{12} \operatorname{tr}_+(1324) \operatorname{tr}_+(1543)^2}{6s_{13} \operatorname{tr}_+(2543)^2} - \frac{s_{45} \operatorname{tr}_+(1234)^2 \operatorname{tr}_+(1534) \operatorname{tr}_+(2453)}{6s_{12}s_{34}s_{35} \operatorname{tr}_+(2543)^2},
 \end{aligned} \tag{4.11}$$

$$\begin{aligned}
 c_{51;23} = & \frac{s_{23}s_{45} \operatorname{tr}_+(1234)^2 \operatorname{tr}_+(1534)}{6s_{12}s_{34} \operatorname{tr}_+(2543)^2} + \frac{2s_{12}s_{23}s_{34}s_{45} \operatorname{tr}_+(1543)}{3 \operatorname{tr}_+(2543)^2} \\
 & - \frac{5s_{23}s_{45} \operatorname{tr}_+(1234) \operatorname{tr}_+(1543)}{2 \operatorname{tr}_+(2543)^2},
 \end{aligned} \tag{4.12}$$

$$c_{\text{rat}} = -\frac{5 \operatorname{tr}_+(1234) \operatorname{tr}_+(1543)}{4s_{34} \operatorname{tr}_+(2543)} + \frac{5s_{23} \operatorname{tr}_+(1243) \operatorname{tr}_+(1543)^2}{2s_{15}s_{34} \operatorname{tr}_+(2543)^2}. \tag{4.13}$$

While for the $(d_s - 2)^2$ amplitude the coefficients are:

$$d_{34}^{(3)} = -\frac{s_{12}s_{15} \operatorname{tr}_+(2543)^2}{12s_{34}}, \tag{4.14a}$$

$$d_{34}^{(2)} = \frac{1}{6}s_{12}s_{15} \operatorname{tr}_+(2543), \tag{4.14b}$$

$$d_{34;51}^{(3)} = -\frac{1}{18}s_{15}s_{23} \operatorname{tr}_+(2543), \tag{4.14c}$$

$$d_{34;51}^{(2)} = -\frac{s_{15} \operatorname{tr}_+(2543)^2}{36s_{12}s_{34}} - \frac{1}{6}s_{15} \operatorname{tr}_+(2543), \tag{4.14d}$$

$$d_{51;23}^{(3)} = \frac{s_{15}s_{23}s_{34}^2s_{45}^2 \operatorname{tr}_+(1245)}{18 \operatorname{tr}_+(2543)^2}, \tag{4.15a}$$

$$d_{51;23}^{(2)} = \frac{s_{12}s_{15}s_{23}s_{34}^2s_{45}^2}{12 \operatorname{tr}_+(2543)^2} - \frac{s_{15}s_{23}s_{34}s_{45}^2 \operatorname{tr}_+(1234)}{6 \operatorname{tr}_+(2543)^2}, \tag{4.15b}$$

$$\begin{aligned}
 d_{\text{rat}} = & \frac{s_{34}}{72} + \frac{5s_{45}}{36} - \frac{\operatorname{tr}_+(1234) \operatorname{tr}_+(1453)}{72s_{14} \operatorname{tr}_+(2543)} - \frac{s_{45} \operatorname{tr}_+(1543)}{72 \operatorname{tr}_+(2543)} + \frac{\operatorname{tr}_+(2543)^2}{72s_{34}^3} \\
 & - \frac{5 \operatorname{tr}_+(1543)^2}{72s_{34} \operatorname{tr}_+(2543)} - \frac{s_{12} \operatorname{tr}_+(1543)^2}{72 \operatorname{tr}_+(2543)^2} + \frac{s_{23}s_{34}s_{45} \operatorname{tr}_+(1543)}{18 \operatorname{tr}_+(2543)^2}.
 \end{aligned} \tag{4.16}$$

These results can also be found in the supplementary material attached to this article.

5 Conclusions

In this article we have presented a new analytic result for a two-loop five-gluon scattering amplitude in QCD. We were able to find a compact representation for the finite remainder of the single-minus helicity configuration after removing the universal infrared poles. We set up a complete tool-chain from generalised unitarity cuts to the coefficients of a basis of pentagon functions for the finite remainder. This numerical algorithm was then evaluated multiple times with finite field arithmetic and the analytic result reconstructed, avoiding the usual large intermediate expressions.

This single-minus amplitude has turned out to be significantly more difficult to compute than the highly symmetric all-plus helicity amplitude that has been known for some time [27, 59]. At the level of the master integrals the single-minus amplitude was of similar complexity to the maximal-helicity-violating (MHV) configurations. However, after removing the contribution from the universal poles, the finite remainder was simple and contained only up to weight two polylogarithms. This makes the final answer simpler than the more general MHV case which will have up to weight four polylogarithms.

Nevertheless, the techniques presented here are not helicity dependent so we hope to find applications to the remaining independent planar helicity amplitudes in the near future. The last few months have also seen progress on non-planar integrals [62, 64, 65] for five-point scattering which is encouraging for applications to non-planar amplitudes.

Acknowledgments

We are extremely grateful to Johannes Henn, Thomas Gehrmann, Adriano Lo Presti, Fabrizio Caola and Calum Milloy for useful discussions. SB is supported by an STFC Rutherford Fellowship ST/L004925/1 and CBH and HBH are supported by Rutherford Grant ST/M004104/1. This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 772099. This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement 746223.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] J.R. Andersen et al., *Les Houches 2017: Physics at TeV Colliders Standard Model Working Group Report*, in *10th Les Houches Workshop on Physics at TeV Colliders (PhysTeV 2017)*, Les Houches, France, June 5–23, 2017 (2018) [[arXiv:1803.07977](https://arxiv.org/abs/1803.07977)].
- [2] G. Ossola, C.G. Papadopoulos and R. Pittau, *Reducing full one-loop amplitudes to scalar integrals at the integrand level*, *Nucl. Phys. B* **763** (2007) 147 [[hep-ph/0609007](https://arxiv.org/abs/hep-ph/0609007)] [[INSPIRE](https://inspirehep.net/literature/570489)].
- [3] G. Ossola, C.G. Papadopoulos and R. Pittau, *CutTools: A Program implementing the OPP reduction method to compute one-loop amplitudes*, *JHEP* **03** (2008) 042 [[arXiv:0711.3596](https://arxiv.org/abs/0711.3596)] [[INSPIRE](https://inspirehep.net/literature/152668)].

- [4] P. Mastrolia, G. Ossola, C.G. Papadopoulos and R. Pittau, *Optimizing the Reduction of One-Loop Amplitudes*, *JHEP* **06** (2008) 030 [[arXiv:0803.3964](#)] [[INSPIRE](#)].
- [5] P. Mastrolia, G. Ossola, T. Reiter and F. Tramontano, *Scattering AMplitudes from Unitarity-based Reduction Algorithm at the Integrand-level*, *JHEP* **08** (2010) 080 [[arXiv:1006.0710](#)] [[INSPIRE](#)].
- [6] G. Cullen et al., *Automated One-Loop Calculations with GoSam*, *Eur. Phys. J. C* **72** (2012) 1889 [[arXiv:1111.2034](#)] [[INSPIRE](#)].
- [7] Z. Bern, L.J. Dixon, D.C. Dunbar and D.A. Kosower, *One loop n point gauge theory amplitudes, unitarity and collinear limits*, *Nucl. Phys. B* **425** (1994) 217 [[hep-ph/9403226](#)] [[INSPIRE](#)].
- [8] Z. Bern, L.J. Dixon, D.C. Dunbar and D.A. Kosower, *Fusing gauge theory tree amplitudes into loop amplitudes*, *Nucl. Phys. B* **435** (1995) 59 [[hep-ph/9409265](#)] [[INSPIRE](#)].
- [9] Z. Bern, L.J. Dixon and D.A. Kosower, *One loop amplitudes for e^+e^- to four partons*, *Nucl. Phys. B* **513** (1998) 3 [[hep-ph/9708239](#)] [[INSPIRE](#)].
- [10] R. Britto, F. Cachazo and B. Feng, *Generalized unitarity and one-loop amplitudes in $N = 4$ super-Yang-Mills*, *Nucl. Phys. B* **725** (2005) 275 [[hep-th/0412103](#)] [[INSPIRE](#)].
- [11] D. Forde, *Direct extraction of one-loop integral coefficients*, *Phys. Rev. D* **75** (2007) 125019 [[arXiv:0704.1835](#)] [[INSPIRE](#)].
- [12] R.K. Ellis, W.T. Giele and Z. Kunszt, *A Numerical Unitarity Formalism for Evaluating One-Loop Amplitudes*, *JHEP* **03** (2008) 003 [[arXiv:0708.2398](#)] [[INSPIRE](#)].
- [13] W.T. Giele, Z. Kunszt and K. Melnikov, *Full one-loop amplitudes from tree amplitudes*, *JHEP* **04** (2008) 049 [[arXiv:0801.2237](#)] [[INSPIRE](#)].
- [14] C.F. Berger et al., *An Automated Implementation of On-Shell Methods for One-Loop Amplitudes*, *Phys. Rev. D* **78** (2008) 036003 [[arXiv:0803.4180](#)] [[INSPIRE](#)].
- [15] P. Mastrolia and G. Ossola, *On the Integrand-Reduction Method for Two-Loop Scattering Amplitudes*, *JHEP* **11** (2011) 014 [[arXiv:1107.6041](#)] [[INSPIRE](#)].
- [16] D.A. Kosower and K.J. Larsen, *Maximal Unitarity at Two Loops*, *Phys. Rev. D* **85** (2012) 045017 [[arXiv:1108.1180](#)] [[INSPIRE](#)].
- [17] S. Badger, H. Frellesvig and Y. Zhang, *Hepta-Cuts of Two-Loop Scattering Amplitudes*, *JHEP* **04** (2012) 055 [[arXiv:1202.2019](#)] [[INSPIRE](#)].
- [18] Y. Zhang, *Integrand-Level Reduction of Loop Amplitudes by Computational Algebraic Geometry Methods*, *JHEP* **09** (2012) 042 [[arXiv:1205.5707](#)] [[INSPIRE](#)].
- [19] S. Badger, H. Frellesvig and Y. Zhang, *An Integrand Reconstruction Method for Three-Loop Amplitudes*, *JHEP* **08** (2012) 065 [[arXiv:1207.2976](#)] [[INSPIRE](#)].
- [20] P. Mastrolia, E. Mirabella, G. Ossola and T. Peraro, *Scattering Amplitudes from Multivariate Polynomial Division*, *Phys. Lett. B* **718** (2012) 173 [[arXiv:1205.7087](#)] [[INSPIRE](#)].
- [21] P. Mastrolia, E. Mirabella, G. Ossola and T. Peraro, *Integrand-Reduction for Two-Loop Scattering Amplitudes through Multivariate Polynomial Division*, *Phys. Rev. D* **87** (2013) 085026 [[arXiv:1209.4319](#)] [[INSPIRE](#)].
- [22] P. Mastrolia, E. Mirabella, G. Ossola and T. Peraro, *Multiloop Integrand Reduction for Dimensionally Regulated Amplitudes*, *Phys. Lett. B* **727** (2013) 532 [[arXiv:1307.5832](#)] [[INSPIRE](#)].

- [23] S. Caron-Huot and K.J. Larsen, *Uniqueness of two-loop master contours*, *JHEP* **10** (2012) 026 [[arXiv:1205.0801](#)] [[INSPIRE](#)].
- [24] P. Mastrolia, T. Peraro and A. Primo, *Adaptive Integrand Decomposition in parallel and orthogonal space*, *JHEP* **08** (2016) 164 [[arXiv:1605.03157](#)] [[INSPIRE](#)].
- [25] T. Peraro, *Scattering amplitudes over finite fields and multivariate functional reconstruction*, *JHEP* **12** (2016) 030 [[arXiv:1608.01902](#)] [[INSPIRE](#)].
- [26] S. Abreu, F. Febres Cordero, H. Ita, M. Jaquier and B. Page, *Subleading Poles in the Numerical Unitarity Method at Two Loops*, *Phys. Rev. D* **95** (2017) 096011 [[arXiv:1703.05255](#)] [[INSPIRE](#)].
- [27] S. Badger, H. Frellesvig and Y. Zhang, *A Two-Loop Five-Gluon Helicity Amplitude in QCD*, *JHEP* **12** (2013) 045 [[arXiv:1310.1051](#)] [[INSPIRE](#)].
- [28] S. Badger, G. Mogull, A. Ochirov and D. O’Connell, *A Complete Two-Loop, Five-Gluon Helicity Amplitude in Yang-Mills Theory*, *JHEP* **10** (2015) 064 [[arXiv:1507.08797](#)] [[INSPIRE](#)].
- [29] D.C. Dunbar and W.B. Perkins, *Two-loop five-point all plus helicity Yang-Mills amplitude*, *Phys. Rev. D* **93** (2016) 085029 [[arXiv:1603.07514](#)] [[INSPIRE](#)].
- [30] D.C. Dunbar, G.R. Jehu and W.B. Perkins, *The two-loop n-point all-plus helicity amplitude*, *Phys. Rev. D* **93** (2016) 125006 [[arXiv:1604.06631](#)] [[INSPIRE](#)].
- [31] D.C. Dunbar, G.R. Jehu and W.B. Perkins, *Two-loop six gluon all plus helicity amplitude*, *Phys. Rev. Lett.* **117** (2016) 061602 [[arXiv:1605.06351](#)] [[INSPIRE](#)].
- [32] S. Badger, G. Mogull and T. Peraro, *Local integrands for two-loop all-plus Yang-Mills amplitudes*, *JHEP* **08** (2016) 063 [[arXiv:1606.02244](#)] [[INSPIRE](#)].
- [33] D.C. Dunbar, J.H. Godwin, G.R. Jehu and W.B. Perkins, *Analytic all-plus-helicity gluon amplitudes in QCD*, *Phys. Rev. D* **96** (2017) 116013 [[arXiv:1710.10071](#)] [[INSPIRE](#)].
- [34] S. Badger, C. Brønnum-Hansen, H.B. Hartanto and T. Peraro, *First look at two-loop five-gluon scattering in QCD*, *Phys. Rev. Lett.* **120** (2018) 092001 [[arXiv:1712.02229](#)] [[INSPIRE](#)].
- [35] S. Abreu, F. Febres Cordero, H. Ita, B. Page and M. Zeng, *Planar Two-Loop Five-Gluon Amplitudes from Numerical Unitarity*, *Phys. Rev. D* **97** (2018) 116014 [[arXiv:1712.03946](#)] [[INSPIRE](#)].
- [36] S. Badger et al., *Applications of integrand reduction to two-loop five-point scattering amplitudes in QCD*, *PoS(LL2018)006* (2018) [[arXiv:1807.09709](#)] [[INSPIRE](#)].
- [37] S. Abreu, F. Febres Cordero, H. Ita, B. Page and V. Sotnikov, *Planar Two-Loop Five-Parton Amplitudes from Numerical Unitarity*, *JHEP* **11** (2018) 116 [[arXiv:1809.09067](#)] [[INSPIRE](#)].
- [38] R.H. Boels, Q. Jin and H. Lüo, *Efficient integrand reduction for particles with spin*, [arXiv:1802.06761](#) [[INSPIRE](#)].
- [39] H.A. Chawdhry, M.A. Lim and A. Mitov, *Two-loop five-point massless QCD amplitudes within the IBP approach*, [arXiv:1805.09182](#) [[INSPIRE](#)].
- [40] F.V. Tkachov, *A Theorem on Analytical Calculability of Four Loop Renormalization Group Functions*, *Phys. Lett. B* **100** (1981) 65 [[INSPIRE](#)].
- [41] K.G. Chetyrkin and F.V. Tkachov, *Integration by Parts: The Algorithm to Calculate β -functions in 4 Loops*, *Nucl. Phys. B* **192** (1981) 159 [[INSPIRE](#)].

- [42] J. Gluza, K. Kajda and D.A. Kosower, *Towards a Basis for Planar Two-Loop Integrals*, *Phys. Rev. D* **83** (2011) 045012 [[arXiv:1009.0472](#)] [[INSPIRE](#)].
- [43] H. Ita, *Two-loop Integrand Decomposition into Master Integrals and Surface Terms*, *Phys. Rev. D* **94** (2016) 116015 [[arXiv:1510.05626](#)] [[INSPIRE](#)].
- [44] K.J. Larsen and Y. Zhang, *Integration-by-parts reductions from unitarity cuts and algebraic geometry*, *Phys. Rev. D* **93** (2016) 041701 [[arXiv:1511.01071](#)] [[INSPIRE](#)].
- [45] A. Georgoudis, K.J. Larsen and Y. Zhang, *Azurite: An algebraic geometry based package for finding bases of loop integrals*, *Comput. Phys. Commun.* **221** (2017) 203 [[arXiv:1612.04252](#)] [[INSPIRE](#)].
- [46] D.A. Kosower, *Direct Solution of Integration-by-Parts Systems*, *Phys. Rev. D* **98** (2018) 025008 [[arXiv:1804.00131](#)] [[INSPIRE](#)].
- [47] J. Böhm, A. Georgoudis, K.J. Larsen, H. Schönemann and Y. Zhang, *Complete integration-by-parts reductions of the non-planar hexagon-box via module intersections*, *JHEP* **09** (2018) 024 [[arXiv:1805.01873](#)] [[INSPIRE](#)].
- [48] J. Böhm, A. Georgoudis, K.J. Larsen, M. Schulze and Y. Zhang, *Complete sets of logarithmic vector fields for integration-by-parts identities of Feynman integrals*, *Phys. Rev. D* **98** (2018) 025023 [[arXiv:1712.09737](#)] [[INSPIRE](#)].
- [49] S. Laporta, *High precision calculation of multiloop Feynman integrals by difference equations*, *Int. J. Mod. Phys. A* **15** (2000) 5087 [[hep-ph/0102033](#)] [[INSPIRE](#)].
- [50] A. von Manteuffel and C. Studerus, *Reduze 2 — Distributed Feynman Integral Reduction*, [arXiv:1201.4330](#) [[INSPIRE](#)].
- [51] A.V. Smirnov, *FIRE5: a C++ implementation of Feynman Integral REduction*, *Comput. Phys. Commun.* **189** (2015) 182 [[arXiv:1408.2372](#)] [[INSPIRE](#)].
- [52] P. Maierhöfer, J. Usovitsch and P. Uwer, *Kira — A Feynman integral reduction program*, *Comput. Phys. Commun.* **230** (2018) 99 [[arXiv:1705.05610](#)] [[INSPIRE](#)].
- [53] A. von Manteuffel and R.M. Schabinger, *A novel approach to integration by parts reduction*, *Phys. Lett. B* **744** (2015) 101 [[arXiv:1406.4513](#)] [[INSPIRE](#)].
- [54] A.V. Kotikov, *Differential equations method: New technique for massive Feynman diagrams calculation*, *Phys. Lett. B* **254** (1991) 158 [[INSPIRE](#)].
- [55] T. Gehrmann and E. Remiddi, *Differential equations for two loop four point functions*, *Nucl. Phys. B* **580** (2000) 485 [[hep-ph/9912329](#)] [[INSPIRE](#)].
- [56] J.M. Henn, *Multiloop integrals in dimensional regularization made simple*, *Phys. Rev. Lett.* **110** (2013) 251601 [[arXiv:1304.1806](#)] [[INSPIRE](#)].
- [57] C.G. Papadopoulos, *Simplified differential equations approach for Master Integrals*, *JHEP* **07** (2014) 088 [[arXiv:1401.6057](#)] [[INSPIRE](#)].
- [58] A. von Manteuffel, E. Panzer and R.M. Schabinger, *A quasi-finite basis for multi-loop Feynman integrals*, *JHEP* **02** (2015) 120 [[arXiv:1411.7392](#)] [[INSPIRE](#)].
- [59] T. Gehrmann, J.M. Henn and N.A. Lo Presti, *Analytic form of the two-loop planar five-gluon all-plus-helicity amplitude in QCD*, *Phys. Rev. Lett.* **116** (2016) 062001 [Erratum *ibid.* **116** (2016) 189903] [[arXiv:1511.05409](#)] [[INSPIRE](#)].

- [60] C.G. Papadopoulos, D. Tommasini and C. Wever, *The Pentabox Master Integrals with the Simplified Differential Equations approach*, *JHEP* **04** (2016) 078 [[arXiv:1511.09404](#)] [[INSPIRE](#)].
- [61] M. Zeng, *Differential equations on unitarity cut surfaces*, *JHEP* **06** (2017) 121 [[arXiv:1702.02355](#)] [[INSPIRE](#)].
- [62] D. Chicherin, J. Henn and V. Mitev, *Bootstrapping pentagon functions*, *JHEP* **05** (2018) 164 [[arXiv:1712.09610](#)] [[INSPIRE](#)].
- [63] T. Gehrmann, J.M. Henn and N.A. Lo Presti, *Pentagon functions for massless planar scattering amplitudes*, *JHEP* **10** (2018) 103 [[arXiv:1807.09812](#)] [[INSPIRE](#)].
- [64] S. Abreu, B. Page and M. Zeng, *Differential equations from unitarity cuts: nonplanar hexa-box integrals*, *JHEP* **01** (2019) 006 [[arXiv:1807.11522](#)] [[INSPIRE](#)].
- [65] D. Chicherin, T. Gehrmann, J.M. Henn, N.A. Lo Presti, V. Mitev and P. Wasser, *Analytic result for the nonplanar hexa-box integrals*, [arXiv:1809.06240](#) [[INSPIRE](#)].
- [66] Z. Bern, L.J. Dixon and D.A. Kosower, *One loop corrections to five gluon amplitudes*, *Phys. Rev. Lett.* **70** (1993) 2677 [[hep-ph/9302280](#)] [[INSPIRE](#)].
- [67] S. Caron-Huot, L.J. Dixon, M. von Hippel, A.J. McLeod and G. Papathanasiou, *The Double Pentagonal Integral to All Orders*, *JHEP* **07** (2018) 170 [[arXiv:1806.01361](#)] [[INSPIRE](#)].
- [68] L.J. Dixon, J. Drummond, T. Harrington, A.J. McLeod, G. Papathanasiou and M. Spradlin, *Heptagons from the Steinmann Cluster Bootstrap*, *JHEP* **02** (2017) 137 [[arXiv:1612.08976](#)] [[INSPIRE](#)].
- [69] S. Caron-Huot, L.J. Dixon, A. McLeod and M. von Hippel, *Bootstrapping a Five-Loop Amplitude Using Steinmann Relations*, *Phys. Rev. Lett.* **117** (2016) 241601 [[arXiv:1609.00669](#)] [[INSPIRE](#)].
- [70] L.J. Dixon, M. von Hippel and A.J. McLeod, *The four-loop six-gluon NMHV ratio function*, *JHEP* **01** (2016) 053 [[arXiv:1509.08127](#)] [[INSPIRE](#)].
- [71] Ø. Almhelid, C. Duhr, E. Gardi, A. McLeod and C.D. White, *Bootstrapping the QCD soft anomalous dimension*, *JHEP* **09** (2017) 073 [[arXiv:1706.10162](#)] [[INSPIRE](#)].
- [72] S. Catani, *The Singular behavior of QCD amplitudes at two loop order*, *Phys. Lett. B* **427** (1998) 161 [[hep-ph/9802439](#)] [[INSPIRE](#)].
- [73] T. Becher and M. Neubert, *On the Structure of Infrared Singularities of Gauge-Theory Amplitudes*, *JHEP* **06** (2009) 081 [Erratum *ibid.* **11** (2013) 024] [[arXiv:0903.1126](#)] [[INSPIRE](#)].
- [74] T. Becher and M. Neubert, *Infrared singularities of scattering amplitudes in perturbative QCD*, *Phys. Rev. Lett.* **102** (2009) 162001 [Erratum *ibid.* **111** (2013) 199905] [[arXiv:0901.0722](#)] [[INSPIRE](#)].
- [75] E. Gardi and L. Magnea, *Factorization constraints for soft anomalous dimensions in QCD scattering amplitudes*, *JHEP* **03** (2009) 079 [[arXiv:0901.1091](#)] [[INSPIRE](#)].
- [76] A. Hodges, *Eliminating spurious poles from gauge-theoretic amplitudes*, *JHEP* **05** (2013) 135 [[arXiv:0905.1473](#)] [[INSPIRE](#)].
- [77] N. Arkani-Hamed, J.L. Bourjaily, F. Cachazo, S. Caron-Huot and J. Trnka, *The All-Loop Integrand For Scattering Amplitudes in Planar $N = 4$ SYM*, *JHEP* **01** (2011) 041 [[arXiv:1008.2958](#)] [[INSPIRE](#)].

- [78] N. Arkani-Hamed, J.L. Bourjaily, F. Cachazo and J. Trnka, *Local Integrals for Planar Scattering Amplitudes*, *JHEP* **06** (2012) 125 [[arXiv:1012.6032](#)] [[INSPIRE](#)].
- [79] J.L. Bourjaily, E. Herrmann and J. Trnka, *Prescriptive Unitarity*, *JHEP* **06** (2017) 059 [[arXiv:1704.05460](#)] [[INSPIRE](#)].
- [80] F.A. Berends and W.T. Giele, *Recursive Calculations for Processes with n Gluons*, *Nucl. Phys. B* **306** (1988) 759 [[INSPIRE](#)].
- [81] C. Cheung and D. O'Connell, *Amplitudes and Spinor-Helicity in Six Dimensions*, *JHEP* **07** (2009) 075 [[arXiv:0902.0981](#)] [[INSPIRE](#)].
- [82] R.N. Lee, *Presenting LiteRed: a tool for the Loop InTEgrals REDuction*, [arXiv:1212.2685](#) [[INSPIRE](#)].
- [83] Z. Bern, L.J. Dixon and D.A. Kosower, *One loop corrections to two quark three gluon amplitudes*, *Nucl. Phys. B* **437** (1995) 259 [[hep-ph/9409393](#)] [[INSPIRE](#)].